

Limits of Hypergraphs, Removal and Regularity Lemmas. A Non-standard Approach*

GÁBOR ELEK and BALÁZS SZEGEDY

February 1, 2008

Abstract

We study the integral and measure theory of the ultraproduct of finite sets. As a main application we construct limit objects for hypergraph sequences. We give a new proof for the Hypergraph Removal Lemma and the Hypergraph Regularity Lemma.

Contents

1	Introduction	2
2	Analysis on the ultraproduct of finite measure spaces	3
2.1	Ultraproducts of finite sets	3
2.2	Measureable functions and their integrals	4
2.3	Fubini's Theorem and the Integration Rule	6
2.4	Random Partitions	8
2.5	Independent Complement in Separable σ -algebras	10
2.6	Separable Realization	13
3	Applications for Hypergraphs	16
3.1	Hypergraph homomorphisms and convergence	16
3.2	The Hypergraph Removal Lemma	17
3.3	The Hypergraph Limit Object	19
3.4	The Hypergraph Regularity Lemma	20
4	Appendix on basic measure theory	23

*AMS Subject Classification: Primary 05C99, Secondary 82B99

1 Introduction

The so-called Hypergraph Regularity Lemma (Rödl-Skokan [9], Gowers [2], later generalized by Tao [11]) is one of the most exciting result in modern combinatorics. It exists in many different forms, strength and generality. The main message in all of them is that every k -uniform hypergraph can be approximated by a structure which consists of boundedly many random-looking (quasi-random) parts for any given error ϵ . Another common feature of these theorems is that they all come with a corresponding counting lemma [8] which describes how to estimate the frequency of a given small hypergraph from the quasi-random approximation of a large hypergraph. One of the most important applications of this method is that it implies the Hypergraph Removal Lemma (first proved by Nagle, Rödl and Schacht [8]) and by an observation of Solymosi [10] it also implies Szemerédi's celebrated theorem on arithmetic progressions in dense subsets of the integers even in a multidimensional setting.

In this paper we present an analytic approach to the subject. First, for any given sequence of hypergraphs we associate the so-called ultralimit hypergraph, which is a measurable hypergraph in a large (non-separable) probability measure space. The ultralimit method enables us to convert theorems of finite combinatorics to measure theoretic statements on our ultralimit space. In the second step, using separable approximations we translates these measure-theoretic theorems to well-known results on the more familiar Lebesgue spaces. This way in two steps we prove the Hypergraph Removal Lemma from the Lebesgue Density Theorem and the Hypergraph Regularity Lemma from the Rectangular Approximation Lemma of Lebesgue Spaces.

We also construct a Hypergraph Limit Object to convergent hypergraph sequences directly from the ultralimit hypergraph. This construction is the generalization of the limit graph method [1],[6] where limits of sequences of dense graphs are studied. According to a definition by Borgs et. al. [1] a graph sequence is called convergent if the density of any fixed graph in the terms of the sequence is convergent. In a paper by Lovasz and Szegedy [6] it is shown that a convergent graph sequence has a natural limit object which is a two variable function $w : [0, 1]^2 \rightarrow [0, 1]$ with $w(x, y) = w(y, x)$. Informally speaking, w is an infinite analogue of the adjacency matrix. Our main theorem is a generalization of this theorem to k -uniform hypergraphs. We also show that limits of k -uniform hypergraphs can be represented by $2^k - 2$ variable measurable functions $w : [0, 1]^{2^k - 2} \rightarrow [0, 1]$ such that the coordinates are indexed by the proper non empty subsets of $\{1, 2, \dots, k\}$ and w is invariant under the induced action of S_k on the coordinates.

Acknowledgement: We are very indebted to Terence Tao and László Lovász for helpful discussions.

2 Analysis on the ultraproduct of finite measure spaces

2.1 Ultraproducts of finite sets

First we recall the ultraproduct construction of finite probability measure spaces (see [5]). Let $\{X_i\}_{i=1}^\infty$ be finite sets. We always suppose that $|X_1| < |X_2| < |X_3| < \dots$. Let ω be a nonprincipal ultrafilter and $\lim_\omega : l^\infty(\mathbb{N}) \rightarrow \mathbb{R}$ be the corresponding ultralimit. Recall that $\lim_\omega$ is a bounded linear functional such that for any $\epsilon > 0$ and $\{a_n\}_{n=1}^\infty \in l^\infty(\mathbb{N})$

$$\{i \in \mathbb{N} \mid a_i \in [\lim_\omega a_n - \epsilon, \lim_\omega a_n + \epsilon]\} \in \omega.$$

The ultraproduct of the sets X_i is defined as follows.

Let $\tilde{X} = \prod_{i=1}^\infty X_i$. We say that $\tilde{p} = \{p_i\}_{i=1}^\infty, \tilde{q} = \{q_i\}_{i=1}^\infty \in \tilde{X}$ are equivalent, $\tilde{p} \sim \tilde{q}$, if

$$\{i \in \mathbb{N} \mid p_i = q_i\} \in \omega.$$

Define $\mathbf{X} := \tilde{X} / \sim$. Now let $\mathcal{P}(X_i)$ denote the Boolean-algebra of subsets of X_i , with the normalized measure $\mu_i(A) = \frac{|A|}{|X_i|}$. Then let $\tilde{\mathcal{P}} = \prod_{i=1}^\infty \mathcal{P}(X_i)$ and $\mathcal{P} = \tilde{\mathcal{P}} / I$, where I is the ideal of elements $\{A_i\}_{i=1}^\infty$ such that $\{i \in \mathbb{N} \mid A_i = \emptyset\} \in \omega$. Notice that the elements of $\mathcal{P}$ can be identified with certain subsets of $\mathbf{X}$: If

$$\bar{p} = [\{p_i\}_{i=1}^\infty] \in \mathbf{X} \text{ and } \bar{A} = [\{A_i\}_{i=1}^\infty] \in \mathcal{P}$$

then $\bar{p} \in \bar{A}$ if $\{i \in \mathbb{N} \mid p_i \in A_i\} \in \omega$. Clearly, if $\bar{A} = [\{A_i\}_{i=1}^\infty], \bar{B} = [\{B_i\}_{i=1}^\infty]$ then

- $\bar{A}^c = [\{A_i^c\}_{i=1}^\infty],$
- $\bar{A} \cup \bar{B} = [\{A_i \cup B_i\}_{i=1}^\infty],$
- $\bar{A} \cap \bar{B} = [\{A_i \cap B_i\}_{i=1}^\infty].$

That is $\mathcal{P}$ is a Boolean algebra on $\mathbf{X}$. Now let $\mu(\bar{A}) = \lim_\omega \mu_i(A_i)$. Then $\mu : \mathcal{P} \rightarrow \mathbb{R}$ is a finitely additive probability measure.

Definition 2.1 $N \subseteq \mathbf{X}$ is a **nullset** if for any $\epsilon > 0$ there exists a set $\bar{A}_\epsilon \in \mathcal{P}$ such that $N \subseteq \bar{A}_\epsilon$ and $\mu(\bar{A}_\epsilon) \leq \epsilon$. The set of nullsets is denoted by $\mathcal{N}$.

Proposition 2.1 $\mathcal{N}$ satisfies the following properties:

- if $N \in \mathcal{N}$ and $M \subseteq N$, then $M \in \mathcal{N}$.
- If $\{N_k\}_{k=1}^\infty$ are elements of $\mathcal{N}$ then $\cup_{k=1}^\infty N_k \in \mathcal{N}$ as well.

Proof. We need the following lemma.

Lemma 2.1 If $\{\bar{A}_k\}_{k=1}^\infty$ are elements of $\mathcal{P}$ and $\lim_{l \rightarrow \infty} \mu(\cup_{k=1}^l \bar{A}_k) = t$ then there exists an element $\bar{B} \in \mathcal{P}$ such that $\mu(\bar{B}) = t$ and $\bar{A}_k \subseteq \bar{B}$ for all $k \in \mathbb{N}$.

Proof. Let $\overline{B}_l = \cup_{k=1}^l \overline{A}_k$, $\mu(\overline{B}_l) = t_l$, $\lim_{l \rightarrow \infty} t_l = t$. Let

$$T_l = \left\{ i \in \mathbb{N} \mid |\mu_i(\cup_{k=1}^l A_k^i) - t_l| \leq \frac{1}{2^l} \right\},$$

where $\overline{A}_k = [\{A_k^i\}_{i=1}^\infty]$. Observe that $T_l \in \omega$. If $i \in \cap_{l=1}^m T_l$ but $i \notin T_{m+1}$, then let $C_i = \cup_{k=1}^m A_k^i$. If $i \in T_l$ for all $l \in \mathbb{N}$, then clearly $\mu_i(\cup_{k=1}^\infty A_k^i) = t$ and we set $C_i := \cup_{k=1}^\infty A_k^i$. Let $\overline{B} := [\{C_i\}_{i=1}^\infty]$. Then $\mu(\overline{B}) = t$ and for any $k \in \mathbb{N}$: $\overline{A}_k \subseteq \overline{B}$. ■

Now suppose that for any $j \geq 1$, $\overline{A}_j \in \mathcal{N}$. Let $\overline{B}_j^\epsilon \in \mathcal{P}$ such that $\overline{A}_j \subseteq \overline{B}_j^\epsilon$ and $\mu(\overline{B}_j^\epsilon) < \epsilon \frac{1}{2^j}$. Then by the previous lemma, there exists $\overline{B}^\epsilon \in \mathcal{P}$ such that for any $j \geq 1$ $\overline{B}_j^\epsilon \subseteq \overline{B}^\epsilon$ and $\mu(\overline{B}^\epsilon) \leq \epsilon$. Since $\cup_{j=1}^\infty \overline{A}_j \subseteq \overline{B}^\epsilon$, our proposition follows. ■

Definition 2.2 We call $B \subseteq \mathbf{X}$ a **measureable set** if there exists $\tilde{B} \in \mathcal{P}$ such that $B \Delta \tilde{B} \in \mathcal{N}$.

Theorem 1 The measurable sets form a σ -algebra $\mathcal{B}_\omega$ and $\mu(B) = \mu(\tilde{B})$ defines a probability measure on $\mathcal{B}_\omega$.

Proof. We call two measurable sets B and B' equivalent, $B \cong B'$ if $B \Delta B' \in \mathcal{N}$. Clearly, if $A \cong A'$, $B \cong B'$ then $A^c \cong (A')^c$, $A \cup B \cong A' \cup B'$, $A \cap B \cong A' \cap B'$. Also if $A, B \in \mathcal{P}$ and $A \cong B$, then $\mu(A) = \mu(B)$. That is the measurable sets form a Boolean algebra with a finitely additive measure. Hence it is enough to prove that if $\overline{A}_k \in \mathcal{P}$ are disjoint sets, then there exists $\overline{A} \in \mathcal{P}$ such that $\cup_{k=1}^\infty \overline{A}_k \cong \overline{A}$ and $\mu(\overline{A}) = \sum_{k=1}^\infty \mu(\overline{A}_k)$. Note that by Lemma 2.1 there exists $\overline{A} \in \mathcal{P}$ such that $\mu(\overline{A}) = \sum_{k=1}^\infty \mu(\overline{A}_k)$ and $\overline{A}_k \subseteq \overline{A}$ for all $k \geq 1$. Then for any $j \geq 1$,

$$\overline{A} \setminus \cup_{k=1}^\infty \overline{A}_k \subseteq \overline{A} \setminus \cup_{k=1}^j \overline{A}_k \in \mathcal{P}.$$

Since $\lim_{j \rightarrow \infty} \mu(\overline{A} \setminus \cup_{k=1}^j \overline{A}_k) = 0$, $\overline{A} \setminus \cup_{k=1}^\infty \overline{A}_k \in \mathcal{N}$ thus $\cup_{k=1}^\infty \overline{A}_k \cong \overline{A}$. ■

Hence we constructed an atomless probability measure space $(\mathbf{X}, \mathcal{B}_\omega, \mu)$. Note that this space is non-separable, that is it is not measurably equivalent to the interval with the Lebesgue measure.

2.2 Measureable functions and their integrals

Let $\{X_i\}_{i=1}^\infty$ be finite sets as in the previous section and $f_i : X_i \rightarrow [-d, d]$ be real functions, where $d > 0$. Then one can define a function $f : \mathbf{X} \rightarrow [-d, d]$ whose value at $\bar{p} = [\{p_i\}_{i=1}^\infty]$ is the ultralimit of $\{f_i(p_i)\}_{i=1}^\infty$. We say that f is the ultralimit of the functions $\{f_i\}_{i=1}^\infty$. From now on we call such bounded functions **ultralimit functions**.

Lemma 2.2 *The ultralimit functions are measurable on $\mathbf{X}$ and*

$$\int_{\mathbf{X}} f d\mu = \lim_{\omega} \frac{\sum_{p \in X_i} f_i(p)}{|X_i|}.$$

Proof. Let $-d \leq a \leq b \leq d$ be real numbers. It is enough to prove that $f_{[a,b]} = \{\bar{p} \in \mathbf{X} \mid a \leq f(\bar{p}) \leq b\}$ is measurable. Let $f_{[a,b]}^i = \{p \in X_i \mid a \leq f_i(p) \leq b\}$. Note that $[\{f_{[a,b]}^i\}_{i=1}^\infty]$ is not necessarily equal to $f_{[a,b]}$. Nevertheless if

$$P_n := [\{f_{[a-\frac{1}{n}, b+\frac{1}{n}]}^i\}_{i=1}^\infty],$$

then $P_n \in \mathcal{P}$ and $f_{[a,b]} = \cap_{n=1}^\infty P_n$. This shows that $f_{[a,b]}$ is a measurable set. Hence the function f is measurable. Now we prove the integral formula. Let us consider the function g_i on X_i which takes the value $\frac{j}{2^k}$ if f_i takes a value not greater than $\frac{j}{2^k}$ but less than $\frac{j+1}{2^k}$ for $-N_k \leq j \leq N_k$, where $N_k = [d2^k] + 1$. Clearly $|\lim_{\omega} g_i - f| \leq \frac{1}{2^k}$ on $\mathbf{X}$. Observe that $g = \lim_{\omega} g_i$ is a measurable step-function on $\mathbf{X}$ taking the value $\frac{j}{2^k}$ on $C_j = [\{f_{[\frac{j}{2^k}, \frac{j+1}{2^k}]}^i\}_{i=1}^\infty]$. Hence,

$$\int_X g d\mu = \lim_{\omega} \left(\sum_{j=-N_k}^{N_k} \frac{|f_{[\frac{j}{2^k}, \frac{j+1}{2^k}]}^i|}{|X_i|} \frac{j}{2^k} \right).$$

Also, $|g - f| \leq \frac{1}{2^k}$ on $\mathbf{X}$ uniformly, that is $|\int_{\mathbf{X}} f d\mu - \int_{\mathbf{X}} g d\mu| \leq \frac{1}{2^k}$. Notice that for any $i \geq 1$

$$\left| \sum_{j=-N_k}^{N_k} \frac{|f_{[\frac{j}{2^k}, \frac{j+1}{2^k}]}^i|}{|X_i|} \frac{j}{2^k} - \frac{\sum_{p \in X_i} f_i(p)}{|X_i|} \right| \leq \frac{1}{2^k}.$$

Therefore for each $k \geq 1$,

$$\left| \int_{\mathbf{X}} f d\mu - \lim_{\omega} \frac{\sum_{p \in X_i} f_i(p)}{|X_i|} \right| \leq \frac{1}{2^{k-1}}.$$

Thus our lemma follows. $\blacksquare$

Theorem 2 *For every measurable function $f : \mathbf{X} \rightarrow [-d, d]$, there exists a sequence of functions $f_i : X_i \rightarrow [-d, d]$ such that the ultralimit of the sequence $\{f_i\}_{i=1}^\infty$ is almost everywhere equals to f . That is any element of $L^\infty(\mathbf{X}, \mathcal{B}_\omega, \mu)$ can be represented by an ultralimit function.*

Proof. Recall a standard result of measure theory. If f is a bounded measurable function on $\mathbf{X}$, then there exists a sequence of bounded stepfunctions $\{h_k\}_{k=1}^\infty$ such that

- $f = \sum_{k=1}^\infty h_k$
- $|h_k| \leq \frac{1}{2^{k-1}}$, if $k > 1$.
- $h_k = \sum_{n=1}^{n_k} c_n^k \chi_{A_n^k}$, where $\cup_{n=1}^{n_k} A_n^k = \mathbf{X}$ is a measurable partition, $c_n^k \in \mathbb{R}$ if $1 \leq n \leq n_k$.

Now let $B_n^k \in \mathcal{P}$ such that $\mu(A_n^k \triangle B_n^k) = 0$. We can suppose that $\cup_{n=1}^{n_k} B_n^k$ is a partition of $\mathbf{X}$. Let $h'_k = \sum_{n=1}^{n_k} c_n^k \chi_{B_n^k}$ and $f' = \sum_{k=1}^{\infty} h'_k$. Then clearly $f' = f$ almost everywhere. We show that f' is an ultralimit function.

Let $B_n^k = [\{B_{n,i}^k\}_{i=1}^{\infty}]$. We set $T_k \subset \mathbb{N}$ as the set of integers i for which $\cup_{n=1}^{n_k} B_{n,i}^k$ is a partition of X_i . Then obviously, $T_k \in \omega$. Now we use our diagonalizing trick again. If $i \notin T_1$ let $s_i \equiv 0$. If $i \in T_1, i \in T_2, \dots, i \in T_k, i \notin T_{k+1}$ then define $s_i := \sum_{j=1}^k (\sum_{n=1}^{n_j} c_n^j \chi_{B_{n,i}^j})$. If $i \in T_k$ for each $k \geq 1$ then set $s_i := \sum_{j=1}^i (\sum_{n=1}^{n_j} c_n^j \chi_{B_{n,i}^j})$. Now let $\bar{p} \in B_{j_1}^1 \cap B_{j_2}^2 \cap \dots \cap B_{j_k}^k$. Then

$$|(\lim_{\omega} s_i)(\bar{p}) - f'(\bar{p})| \leq \frac{1}{2^{k-1}}.$$

Since this inequality holds for each $k \geq 1$, $f' \equiv \lim_{\omega} s_i$. $\blacksquare$

2.3 Fubini's Theorem and the Integration Rule

We fix a natural number k and we denote by $[k]$ the set $\{1, 2, \dots, k\}$. Let $X_{i,1}, X_{i,2}, \dots, X_{i,k}$ be k copies of the finite set X_i and for a subset $A \subseteq \{1, 2, \dots, k\}$ let $X_{i,A}$ denote the direct product $\bigoplus_{j \in A} X_{i,j}$. Let $\mathbf{X}^A$ denote the ultra product of the sets $X_{i,A}$, with a Boolean algebra $\mathcal{P}_A$. There is a natural map $p_A : \mathbf{X}^{[k]} \rightarrow \mathbf{X}^A$ (the projection). Let $\mathcal{B}_A$ be the σ -algebra of measurable subsets in $\mathbf{X}^A$ as defined in the previous sections. Define $\sigma(A)$ as $p_A^{-1}(\mathcal{B}_A)$, the σ -algebra of measurable sets depending only on the A -coordinates together with the probability measure μ_A . For a nonempty subset $A \subseteq [k]$ let A^* denote the set system $\{B | B \subseteq A, |B| = |A| - 1\}$ and let $\sigma(A)^*$ denote the σ -algebra $\langle \sigma(B) | B \in A^* \rangle$. An interesting fact is (as it will turn out in subsection 2.4) that $\sigma(A)^*$ is strictly smaller than $\sigma(A)$.

Lemma 2.3 *Let $A, B \subseteq [k]$ and let $f : \mathbf{X}^{[k]} \rightarrow \mathbb{R}$ be a bounded $\sigma(B)$ -measurable ultralimit function. Then for all $y \in \mathbf{X}^{A^c}$ the function f_y is $\sigma(A \cap B)$ -measurable, where A^c denotes the complement of A in $[k]$ and $f_y(x) = f(x, y)$.*

Proof. Let $f : \mathbf{X}^{[k]} \rightarrow \mathbb{R}$ be a $\sigma(B)$ -measurable ultralimit function. It is easy to see that the finite approximation functions $f_i : X_{i,1} \times X_{i,2} \times \dots \times X_{i,k}$ constructed in Theorem 2 depend only on the B -coordinates, since $\sigma(B)$ -measurable functions can be approximated by $\sigma(B)$ -measurable stepfunctions. Let $y \in \mathbf{X}^{A^c}$, $y = [\{y_i\}_{i=1}^{\infty}]$. Then f_y is the ultralimit of the functions $f_i^{y_i}$. Clearly $f_i^{y_i}$ depends only on the $A \cap B$ -coordinates, thus the ultralimit f_y is $\sigma(A \cap B)$ -measurable. $\blacksquare$

Theorem 3 (Fubini's Theorem) *Let $A \subseteq [k]$ and let $f : \mathbf{X}_k \rightarrow \mathbb{R}$ be a $\sigma([k])$ -measurable*

ultralimit function. Then

$$\int_{\mathbf{X}^{[k]}} f(p) d\mu_{[k]}(p) = \int_{\mathbf{X}^{A^c}} \left(\int_{\mathbf{X}^A} f_y(x) d\mu_A(x) \right) d\mu_{A^c}(y)$$

Proof. Let f be the ultralimit of $\{f_i : X_{i,[k]} \rightarrow \mathbb{R}\}_{i=1}^\infty$. Define the functions $\overline{f}_i : X_{i,A^c} \rightarrow [-d, d]$ by

$$\overline{f}_i(y) = |X_{i,A}|^{-1} \sum_{x \in X_{i,A}} f_i(x, y).$$

By Lemma 2.2

$$\lim_{\omega} \overline{f}_i(y) = \int_{\mathbf{X}^A} f(x, y) d\mu_A(x).$$

Applying Lemma 2.2 again for the functions $\overline{f}_i$, we obtain that

$$\lim_{\omega} |X_{i,A^c}|^{-1} \sum_{y \in X_{i,A^c}} \overline{f}_i(y) = \int_{\mathbf{X}^{A^c}} \left(\int_{\mathbf{X}^A} f(x, y) d\mu_A(x) \right) d\mu_{A^c}(y).$$

This completes the proof, since

$$|X_{i,A^c}|^{-1} \sum_{y \in X_{i,A^c}} \overline{f}_i(y) = \frac{\sum_{p \in X_i} f_i(p)}{|X_i|}.$$

■

Recall that if $\mathcal{B} \subset \mathcal{A}$ are σ -algebras on X with a measure μ and g is an $\mathcal{A}$ -measurable function on X , then $E(g \mid \mathcal{B})$ is the $\mathcal{B}$ -measurable function (unique up to a zero measure perturbation) with the property that

$$\int_Y E(g \mid \mathcal{B}) d\mu = \int_Y g d\mu,$$

for any $Y \in \mathcal{B}$ (see Appendix).

Theorem 4 (Integration Rule) *Let $g_i : \mathbf{X}^{[k]} \rightarrow \mathbb{R}$ be bounded $\sigma(A_i)$ -measurable functions for $i = 1, 2, \dots, m$. Let B denote the sigma algebra generated by $\sigma(A_1 \cap A_2), \sigma(A_1 \cap A_3), \dots, \sigma(A_1 \cap A_m)$. Then*

$$\int_{\mathbf{X}^{[k]}} g_1 g_2 \dots g_m d\mu_{[k]} = \int_{\mathbf{X}^{[k]}} E(g_1 \mid B) g_2 g_3 \dots g_m d\mu_{[k]}.$$

Proof. We may suppose that all g_i are ultralimit functions, since the conditional expectation does not depend on zero measure perturbation. Since g_1 does not depend on the A_1^c coordinates we may suppose that $E(g_1 \mid B)$ does not depend on the A_1^c -coordinates as well. By the previous theorem,

$$\int_{\mathbf{X}^{[k]}} g_1 g_2 g_3 \dots g_m d\mu_{[k]} = \int_{\mathbf{X}^{A_1^c}} \left(\int_{\mathbf{X}^{A_1}} g_1(x) g_2(x, y) \dots g_m(x, y) d\mu_{A_1}(x) \right) d\mu_{A_1^c}(y).$$

Now we obtain by Lemma 2.3 that for all $y \in X_{A_1^c}$ the function

$$x \rightarrow g_2(x, y)g_3(x, y) \dots g_m(x, y) \quad (x \in X_{A_1})$$

is B -measurable. This means that

$$\begin{aligned} & \int_{\mathbf{X}^{A_1}} g_1(x)g_2(x, y) \dots g_m(x, y) d\mu_{A_1}(x) = \\ & = \int_{\mathbf{X}^{A_1}} E(g_1|B)(x)g_2(x, y)g_3(x, y) \dots g_m(x, y) d\mu_{A_1}(x) \end{aligned}$$

for all y in $\mathbf{X}_{A_1^c}$. This completes the proof. $\blacksquare$

Lemma 2.4 (Total Independence) *Let $A_1, A_2, \dots, A_r$ be the list of nonempty subsets of $[k]$, and let $S_1, S_2, \dots, S_r$ be subsets of $\mathbf{X}^{[k]}$ such that $S_i \in \sigma(A_i)$ and $E(S_i|\sigma(A_i)^*)$ is a constant function for every $1 \leq i \leq r$. Then*

$$\mu(S_1 \cap S_2 \cap \dots \cap S_r) = \mu(S_1)\mu(S_2) \dots \mu(S_r).$$

Proof. We can assume that $|A_i| \geq |A_j|$ whenever $j > i$. Let χ_i be the characteristic function of S_i . We have that

$$\mu(S_1 \cap S_2 \cap \dots \cap S_r) = \int_{\mathbf{X}^{[k]}} \chi_1 \chi_2 \dots \chi_r d\mu_{[k]}.$$

The integration rule shows that

$$\begin{aligned} \int_{\mathbf{X}^{[k]}} \chi_i \chi_{i+1} \dots \chi_r d\mu_{[k]} &= \int_{\mathbf{X}^{[k]}} E(\chi_i|\sigma(A_i)^*) \chi_{i+1} \dots \chi_r d\mu_{[k]} \\ &= \mu(S_i) \int_{\mathbf{X}^{[k]}} \chi_{i+1} \chi_{i+2} \dots \chi_r d\mu_{[k]}. \end{aligned}$$

This completes the proof. $\blacksquare$

2.4 Random Partitions

The goal of this section is to prove the following proposition.

Proposition 2.2 *Let $A \subset [k]$ be a subset, then for any $n \geq 1$ there exists a partition $X_A = S_1 \cup S_2 \cup \dots \cup S_n$, such that $E(S_i | \sigma(A)^*) = \frac{1}{n}$.*

Proof. The idea of the proof is that we consider random partitions of X_A and show that by probability one these partitions shall satisfy the property of our proposition. Let $\Omega = \prod_{i=1}^{\infty} \{1, 2, \dots, n\}^{X_{i,A}}$ be the set of $\{1, 2, \dots, n\}$ -valued functions on $\cup_{i=1}^{\infty} X_{i,A}$. Each element f of Ω defines a partition of X_A the following way. Let

$$S_f^{i,j} = \{p \in X_{i,A} \mid f(p) = j\} \quad 1 \leq j \leq n, i \geq 1.$$

$$[\{S_f^{i,j}\}_{i=1}^\infty] = S_f^j.$$

Then $X_A = S_f^1 \cup S_f^2 \cup \dots \cup S_f^n$ is our partition induced by f .

Note that on Ω one has the usual Bernoulli probability measure P ,

$$P(T_{p_1, p_2, \dots, p_r}(i_1, i_2, \dots, i_r)) = \frac{1}{n^r},$$

where

$$T_{p_1, p_2, \dots, p_r}(i_1, i_2, \dots, i_r) = \{f \in \Omega \mid f(p_s) = i_s \ 1 \leq s \leq r\}.$$

A **cylindric intersection set** T in $X_{i,A}$ is a set $T = \cap_{C, C \subsetneq A} T_C$, where $T_C \subset X_{i,C}$. First of all note that the number of different cylindric intersection sets in $X_{i,A}$ is not greater than

$$\prod_{C, C \subsetneq A} 2^{|X_{i,C}|} \leq 2^{(|X_i|^{A-1})2^k}.$$

Let $0 \leq \epsilon \leq \frac{1}{10n}$ be a real number and T be a cylindric intersection set of elements at least $\epsilon|X_{i,A}|$. By the Chernoff-inequality the probability that an $f \in \Omega$ takes the value 1 more than $(\frac{1}{n} + \epsilon)|T|$ -times or less than $(\frac{1}{n} - \epsilon)|T|$ -times on the set T is less than $2 \exp(-c_\epsilon|T|)$, where the positive constant c_ϵ depends only on ϵ . Therefore the probability that there exists a cylindric intersection set $T \subset X_{i,A}$ of size at least $\epsilon|X_{i,A}|$ for which $f \in \Omega$ takes the value 1 more than $(\frac{1}{n} + \epsilon)|T|$ -times or less than $(\frac{1}{n} - \epsilon)|T|$ -times on the set T is less than

$$2^{(|X_i|^{A-1})2^k} 2 \exp(-c_\epsilon \epsilon |X_i|^A).$$

Since $|X_1| < |X_2| < \dots$ by the Borel-Cantelli lemma we have the following lemma.

Lemma 2.5 *For almost all $f \in \Omega$ there exist only finitely many i such that there exists at least one cylindric intersection set $T \subset X_{i,A}$ for which $f \in \Omega$ takes the value 1 more than $(\frac{1}{n} + \epsilon)|T|$ -times or less than $(\frac{1}{n} - \epsilon)|T|$ -times on the set T .*

Now let us consider a cylindric intersection set $Z \subseteq X_A$, $Z = \cap_{C, C \subsetneq A} Z_C$, $Z_C \in X_C$. By the previous lemma, for almost all $f \in \Omega$,

$$\mu(S_f^1 \cap Z) = \frac{1}{n} \mu(Z).$$

Therefore for almost all $f \in \Omega$:

$$\mu(S_f^1 \cap Z') = \frac{1}{n} (\mu(Z')),$$

where Z' is a finite disjoint union of cylindric intersection sets in $\mathbf{X}_A$. Consequently, for almost all $f \in \Omega$,

$$\mu(S_f^1 \cap Y) = \frac{1}{n} (\mu(Y)),$$

where $Y \in \sigma(A)^*$. This shows immediately that $E(S_f^1 \mid \sigma(A)^*) = \frac{1}{n}$ for almost all $f \in \Omega$.

Similarly, $E(S_f^i \mid \sigma(A)^*) = \frac{1}{n}$ for almost all $f \in \Omega$, thus our proposition follows. $\blacksquare$

2.5 Independent Complement in Separable σ -algebras

Let $\mathcal{A}$ be a separable σ -algebra on a set X , and let μ be a probability measure on $\mathcal{A}$. Two sub σ -algebras $\mathcal{B}$ and $\mathcal{C}$ are called independent if $\mu(B \cap C) = \mu(B)\mu(C)$ for every $B \in \mathcal{B}$ and $C \in \mathcal{C}$. We say that $\mathcal{C}$ is an *independent complement* of $\mathcal{B}$ in $\mathcal{A}$ if it is independent from $\mathcal{B}$ and $\langle \mathcal{B}, \mathcal{C} \rangle$ is dense in $\mathcal{A}$.

Definition 2.3 Let $\mathcal{A} \geq \mathcal{B}$ be two σ -algebras on a set X and let μ be a probability measure on $\mathcal{A}$. A $\mathcal{B}$ -random k -partition in $\mathcal{A}$ is a partition $A_1, A_2, \dots, A_k$ of X into $\mathcal{A}$ -measurable sets such that $E(A_i|\mathcal{B}) = 1/k$ for every $i = 1, 2, \dots, k$.

Theorem 5 (Independent Complement) Let $\mathcal{A} \geq \mathcal{B}$ be two separable σ -algebras on a set X and let μ be a probability measure on $\mathcal{A}$. Assume that for every natural number k there exists a $\mathcal{B}$ -random k -partition $\{A_{1,k}, A_{2,k}, \dots, A_{k,k}\}$ in $\mathcal{A}$. Then there is an independent complement $\mathcal{C}$ of $\mathcal{B}$ in $\mathcal{A}$. (Note that this is basically the Maharam-lemma, see [7])

Proof. Let $S_1, S_2, \dots$ be a countable generating system of $\mathcal{A}$ and let $\mathcal{P}_k$ denote the finite Boolean algebra generated by $S_1, S_2, \dots, S_k$ and $\{A_{i,j} | i \leq j \leq k\}$. Let $\mathcal{P}_k^*$ denote the atoms of $\mathcal{P}_k$. It is clear that for every atom $R \in \mathcal{P}_k^*$ we have that $E(R|\mathcal{B}) \leq 1/k$ because R is contained in one of the sets $A_{1,k}, A_{2,k}, \dots, A_{k,k}$. During the proof we fix one $\mathcal{B}$ -measurable version of $E(R|\mathcal{B})$ for every R . The algebra $\mathcal{P}_k$ is a subalgebra of $\mathcal{P}_{k+1}$ for every k and so we can define total orderings on the sets $\mathcal{P}_k^*$ such that if $R_1, R_2 \in \mathcal{P}_k^*$ with $R_1 < R_2$ and $R_3, R_4 \in \mathcal{P}_{k+1}^*$ with $R_3 \subseteq R_1, R_4 \subseteq R_2$ then $R_3 < R_4$. We can assume that $\sum_{R \in \mathcal{P}_k^*} E(R, \mathcal{B})(x) = 1$ for every element in X . It follows that for $k \in \mathbb{N}$, $x \in X$ and $\lambda \in [0, 1)$ there is a unique element $R(x, \lambda, k) \in \mathcal{P}_k^*$ satisfying

$$\sum_{R < R(x, \lambda, k)} E(R|\mathcal{B})(x) \leq \lambda$$

and

$$\sum_{R \leq R(x, \lambda, k)} E(R|\mathcal{B})(x) > \lambda.$$

For an element $R \in \mathcal{P}_k^*$ let $T(R, \lambda, k)$ denote the set of those points $x \in X$ for which $R(x, \lambda, k) = R$. It is easy to see that $T(R, \lambda, k)$ is $\mathcal{B}$ -measurable. Let us define the $\mathcal{A}$ -measurable set $S(\lambda, k)$ by

$$S(\lambda, k) = \bigcup_{R \in \mathcal{P}_k^*} (T(R, \lambda, k) \cap (\cup_{R_2 < R} R_2))$$

and $S'(\lambda, k)$ by

$$S'(\lambda, k) = \bigcup_{R \in \mathcal{P}_k^*} (T(R, \lambda, k) \cap (\cup_{R_2 \leq R} R_2)).$$

Proposition 2.3 (i) $\lambda - \frac{1}{k} \leq E(S(\lambda, k) \mid \mathcal{B})(x) \leq \lambda$ for any $x \in X$.

(ii) If $k < t$, then $S(\lambda, k) \subseteq S(\lambda, t) \subseteq S'(\lambda, k)$.

(iii) $E(S'(\lambda, k) \setminus S(\lambda, k) \mid \mathcal{B})(x) \leq \frac{1}{k}$ for any $x \in X$.

Proof. First observe that

$$\lambda - \frac{1}{k} \leq \sum_{R < R(x, \lambda, k)} E(R \mid \mathcal{B})(x) \leq \lambda,$$

for any $x \in X$. Also, we have

$$S(\lambda, k) = \bigcup_{R, R_1 \in \mathcal{P}_k^*, R < R_1} (R \cap T(R_1, \lambda, k)), \quad S'(\lambda, k) = \bigcup_{R, R_1 \in \mathcal{P}_k^*, R \leq R_1} (R \cap T(R_1, \lambda, k)). \quad (1)$$

That is by the basic property of the conditional expectation:

$$E(S(\lambda, k) \mid \mathcal{B}) = \sum_{R, R_1 \in \mathcal{P}_k^*, R < R_1} E(R \mid \mathcal{B}) \chi_{T(R_1, \lambda, k)}.$$

That is

$$E(S(\lambda, k) \mid \mathcal{B})(x) = \sum_{R < R(x, \lambda, k)} E(R \mid \mathcal{B})(x). \quad (2)$$

and similarly

$$E(S'(\lambda, k) \mid \mathcal{B})(x) = \sum_{R \leq R(x, \lambda, k)} E(R \mid \mathcal{B})(x). \quad (3)$$

Hence (i) and (iii) follows immediately, using the fact that $E(R' \mid \mathcal{B}) \leq \frac{1}{k}$ for any $R' \in \mathcal{P}_k^*$.

Observe that for any $R \in \mathcal{P}_k^*$, $T(R, \lambda, k) = \bigcup_{R' \subseteq R, R' \in \mathcal{P}_t^*} T(R', \lambda, t)$. Hence

$$\begin{aligned} \bigcup_{R, R_1 \in \mathcal{P}_k^*, R < R_1} (R \cap T(R_1, \lambda, k)) &\subseteq \bigcup_{R', R'_1 \in \mathcal{P}_t^*, R' < R'_1} (R' \cap T(R'_1, \lambda, t)) \subseteq \\ &\subseteq \bigcup_{R, R_1 \in \mathcal{P}_k^*, R \leq R_1} (R \cap T(R_1, \lambda, k)) \end{aligned}$$

Thus (1) implies (ii). $\blacksquare$

Lemma 2.6 Let $S(\lambda) = \bigcup_{k=1}^{\infty} S(\lambda, k)$. Then if $\lambda_2 < \lambda_1$, then $S(\lambda_2) < S(\lambda_1)$.

Proof. Note that $x \in S(\lambda_2, k)$ if and only if $x \in R_2$ for some $R_2 < R(x, \lambda_2, k)$. Obviously, $R(x, \lambda_2, k) < R(x, \lambda_1, k)$, thus $x \in S(\lambda_1, k)$. Hence $S(\lambda_2) \subseteq S(\lambda_1)$. $\blacksquare$

Lemma 2.7 $E(S(\lambda) \mid \mathcal{B}) = \lambda$.

Proof. Since $\chi_{S(\lambda, k)} \xrightarrow{L_2(X, \mu)} \chi_{S(\lambda)}$, we have $E(S(\lambda, k) \mid \mathcal{B}) \xrightarrow{L_2(X, \mu)} E(S(\lambda) \mid \mathcal{B})$. That is by (i) of Proposition 2.3 $E(S(\lambda) \mid \mathcal{B}) = \lambda$. $\blacksquare$

The last two lemmas together imply that the sets $S(\lambda)$ generate a σ -algebra $\mathcal{C}$ which is independent from $\mathcal{B}$.

Now we have to show that $\mathcal{B}$ and $\mathcal{C}$ generate $\mathcal{A}$. Let $S \in \mathcal{P}_k$ for some $k \in \mathbb{N}$. We say that S is an interval if there exists an element $R \in \mathcal{P}_k^*$ such that $S = \cup_{R_1 \leq R} R_1$. It is enough to show that any interval $S \in \mathcal{P}_k$ can be generated by $\mathcal{B}$ and $\mathcal{C}$.

Suppose that $\{T_t\}_{t=1}^\infty$ be sets in $\langle \mathcal{B}, \mathcal{C} \rangle$ and $\|E(S | \mathcal{B}) - E(T_t | \mathcal{B})\|^2 \rightarrow 0$. Then $\mu(S \triangle T_t) \rightarrow 0$ as $t \rightarrow \infty$, that is $\mathcal{B}$ and $\mathcal{C}$ generate S . Indeed,

$$\mu(S \triangle T_n)^2 = \|\chi_S - \chi_{T_n}\|^2 \geq \|E(S | \mathcal{B}) - E(T_n | \mathcal{B})\|^2.$$

So let $t \geq k$ be an arbitrary natural number. It is clear that S is an interval in $\mathcal{P}_t$. For a natural number $0 \leq d \leq t-1$ let F_d denote the $\mathcal{B}$ -measurable set on which $E(S | \mathcal{B})$ is in the interval $[\frac{d}{t}, \frac{d+1}{t})$. Now we approximate S by

$$T_t = \bigcup_{d=0}^{t-1} (F_d \cap S(\frac{d}{t})) \in \langle \mathcal{B}, \mathcal{C} \rangle.$$

Lemma 2.8 *For any $x \in X$,*

$$|E(S | \mathcal{B})(x) - E(T_t | \mathcal{B})(x)| \leq \frac{3}{t}.$$

Proof. First note that by Proposition 2.3 (iii)

$$\left| E(S(\frac{d}{t}) | \mathcal{B})(x) - E(S(\frac{d}{t}, t) | \mathcal{B})(x) \right| \leq \frac{1}{t}.$$

Note that

$$E(T_t | \mathcal{B})(x) = \sum_{d=0}^{t-1} \chi_{F_d}(x) E(S(\frac{d}{t}) | \mathcal{B})(x).$$

Suppose that $x \in F_d$. Then

$$\left| E(T_t | \mathcal{B})(x) - \sum_{R' < R(x, \frac{d}{t}, t)} E(R' | \mathcal{B})(x) \right| \leq \frac{1}{t}.$$

On the other hand $E(S | \mathcal{B})(x) = \sum_{R' \leq R} E(R' | \mathcal{B})(x)$ and $\frac{d}{t} \leq \sum_{R' \leq R} E(R' | \mathcal{B})(x) < \frac{d+1}{t}$. That is

$$|E(S | \mathcal{B})(x) - E(T_t | \mathcal{B})(x)| \leq \frac{3}{t}. \quad \blacksquare$$

The Theorem now follows from the Lemma immediately. $\blacksquare$

Definition 2.4 *Let $(X, \mathcal{A}, \mu)$ be a probability space, and assume that a finite group G is acting on X such that $\mathcal{A}$ is G -invariant as a set system. We say that the action of G is free if there is a subset S of X with $\mu(S) = 1/|G|$ such that $S^{g_1} \cap S^{g_2} = \emptyset$ whenever g_1 and g_2 are distinct elements of G .*

We will need the following consequence of Theorem 5.

Lemma 2.9 *Let $\mathcal{A} \geq \mathcal{B}$ be two separable σ -algebras on the set X and let μ be a probability measure on $\mathcal{A}$. Assume that a finite group G is acting on X such that $\mathcal{A}, \mathcal{B}$ and μ are G invariant. Assume furthermore that the action of G on $(X, \mathcal{B}, \mu)$ is free and that there is a $\mathcal{B}$ -random k partition of X in $\mathcal{A}$ for every natural number k . Then there is an independent complement $\mathcal{C}$ in $\mathcal{A}$ for $\mathcal{B}$ such that $\mathcal{C}$ is elementwise G -invariant.*

Proof. Let $S \in \mathcal{B}$ be a set showing that G acts freely on $\mathcal{B}$. Let $\mathcal{A}|_S$ and $\mathcal{B}|_S$ denote the restriction of $\mathcal{A}$ and $\mathcal{B}$ to the set S . It is clear that if $\{A_1, A_2, \dots, A_k\}$ is a $\mathcal{B}$ -random k -partition in $\mathcal{A}$ then $\{S \cap A_1, S \cap A_2, \dots, S \cap A_k\}$ is a $\mathcal{B}|_S$ -random k partition in $\mathcal{A}|_S$. Hence by Theorem 5 there exists an independent complement $\mathcal{C}_1$ of $\mathcal{B}|_S$ in $\mathcal{A}|_S$. The set

$$\mathcal{C} = \left\{ \bigcup_{g \in G} H^g \mid H \in \mathcal{C}_1 \right\}$$

is a σ -algebra because the action of G is free. Note that the elements of $\mathcal{C}$ are G -invariant. Since $E(\bigcup_{g \in G} H^g | \mathcal{B}) = \sum_{g \in G} E(H | \mathcal{B})^g$ we get that the elements of $\mathcal{C}$ are independent from $\mathcal{B}$. It is clear that $\langle \mathcal{C}, \mathcal{B} \rangle$ is dense in $\mathcal{A}$. ■

2.6 Separable Realization

In this section we show how to pass from nonseparable σ -algebras to separable ones.

First note that the symmetric group S_k acts on the space X^k by permuting the coordinates:

$$(x_1, x_2, \dots, x_k)^\pi = (x_{\pi^{-1}(1)}, x_{\pi^{-1}(2)}, \dots, x_{\pi^{-1}(k)}).$$

The group also acts on the subsets of $[k]$ and $\sigma(A)^\pi = \sigma(A^\pi)$, where A^π denotes the image of the subset A under $\pi \in S_k$. We shall denote by S_A the symmetric group acting on the subset S_A .

Definition 2.5 *A separable realization of degree r on $\mathbf{X}^k$, $r \leq k$ is a system of atomless separable σ -algebras $\{l(A) \mid \emptyset \neq A \subseteq [k], |A| \leq r\}$ and functions $\{F_A : \mathbf{X}^k \rightarrow [0, 1] \mid \emptyset \neq A \subseteq [k], |A| \leq r\}$ with the following properties*

1. $l(A)$ is a subset of $\sigma(A)$ and is independent from $\sigma(A)^*$ for every $\emptyset \neq A \subseteq [k]$.
2. $l(A)^\pi = l(A^\pi)$ for every permutation $\pi \in S_k$.
3. $S^\pi = S$ for every $S \in l(A)$ and $\pi \in S_A$.
4. F_A is an $l(A)$ -measurable function which defines a measurable equivalence between the measure algebras of $(\mathbf{X}^k, l(A), \mu^k)$ and $[0, 1]$. (see Appendix)
5. $F_A(x) = F_{A^\pi}(x^\pi)$ for every element $x \in \mathbf{X}^k$, $\pi \in S_k$ and $A \subseteq [k]$.

The main theorem in this section is the following one.

Theorem 6 *For every $\mathbf{H} \in \sigma([k])$ there exists a separable realization of degree k such that $\mathbf{H}$ is measurable in $\langle l(A) \mid \emptyset \neq A \subseteq [k] \rangle$.*

We will need the following three lemmas.

Lemma 2.10 *Let $\mathcal{B} \subseteq \mathcal{A}$ two σ -algebras on a set X , and let μ be a probability measure on $\mathcal{A}$. Then for any separable sub- σ -algebra $\bar{\mathcal{A}}$ of $\mathcal{A}$ there exists a separable sub σ -algebra $\bar{\mathcal{B}}$ of $\mathcal{B}$ such that $E(A|\mathcal{B}) = E(A|\bar{\mathcal{B}})$ for every $A \in \bar{\mathcal{A}}$.*

Proof. We use the fact that $\bar{\mathcal{A}}$ is a separable metric space with the distance $d(A, B) = \mu(A \triangle B)$. Let $W = \{D_1, D_2, \dots\}$ be a countable dense subset of $\bar{\mathcal{A}}$ with the previous distance. Let $C_{p,q}^i = E(D_i \mid \mathcal{B})^{-1}(p, q)$, where $p < q$ are rational numbers. Clearly, $E(D_i \mid \mathcal{B})$ is a $\mathcal{B}_i$ -measurable function, where $\mathcal{B}_i = \langle C_{p,q}^i \mid p < q \in \mathbb{Q} \rangle$. Obviously, $E(D_i \mid \bar{\mathcal{B}}) = E(D_i \mid \mathcal{B})$ for any $i \geq 1$, where $\bar{\mathcal{B}} = \langle \mathcal{B}_i \mid i = 1, 2, \dots \rangle$. Now observe that $E(D_i \mid \bar{\mathcal{B}}) \xrightarrow{L_2} E(D, \bar{\mathcal{B}})$ if $D_i \rightarrow D$. Hence for any $D \in \bar{\mathcal{A}}$, $E(D \mid \bar{\mathcal{B}}) = E(D \mid \mathcal{B})$. ■

Lemma 2.11 *Let $A \subseteq [k]$ be a subset and assume that there are atomless separable σ -algebras $d(\{i\}) \subset \sigma(\{i\})$, $i \in A$ such that $d(\{i\})^\pi = d(\{i^\pi\})$ for every $i \in A$ and $\pi \in S_A$. Then S_A acts freely on $\langle d(\{i\}) \mid i \in A \rangle$.*

Proof. The permutation invariance implies that there is a σ -algebra $\mathcal{A}$ on X such that $P_{\{i\}}^{-1}(\mathcal{A}) = d(\{i\})$ for every $i \in A$. Let $F : \mathbf{X} \rightarrow [0, 1]$ be a $\mathcal{A}$ -measurable measure preserving map. Now we can define the map $G : \mathbf{X}^A \rightarrow [0, 1]^A$ by

$$G(x_{i_1}, x_{i_2}, \dots, x_{i_{|A|}}) := (F(x_{i_1}), F(x_{i_2}), \dots, F(x_{i_{|A|}})).$$

Let us introduce $S' := \{(x_1, x_2, \dots, x_r) \mid x_1 < x_2 < \dots < x_r\} \subset [0, 1]^A$ and $S := G^{-1}(S')$. Clearly $\mu^A(S) = 1/|A|!$ and $S^\pi \cap S^\rho = \emptyset$ for every two different elements $\pi \neq \rho$ in S_A . ■

Lemma 2.12 *Let k be a natural number and assume that for every $A \subseteq [k]$ there is a separable σ -algebra $c(A)$ in $\sigma(A)$. Then for every $A \subseteq [k]$ there is a separable σ -algebra $d(A)$ in $\sigma(A)$ with $c(A) \subseteq d(A)$ such that*

1. $E(R|\langle d(B) \mid B \in A^* \rangle) = E(R|\sigma(A)^*)$ whenever $R \in d(A)$.
2. $d(A)^\pi = d(A^\pi)$ for every element $\pi \in S_k$.
3. $d(B) \subseteq d(A)$ whenever $B \subseteq A$

Proof. First we construct algebras $d'(A)$ recursively. Let $d'([k])$ be $\langle c([k])^\pi \mid \pi \in S_k \rangle$. Assume that we have already constructed the algebras $d'(A)$ for $|A| \geq t$. Let $A \subseteq [k]$ be such that

$|A| = t$. By Lemma 2.10 we can see that there exists a separable subalgebra $d'(A)^*$ of $\sigma(A)^*$ such that $E(R|\sigma(A)^*) = E(R|d'(A)^*)$ for every $R \in d'(A)$. Since $\sigma(A)^*$ is generated by the algebras $\{\sigma(B)|B \in A^*\}$ we have that every element of $\sigma(A)^*$ is a countable expression of some sets in these algebras. This implies that any separable sub σ -algebra of $\sigma(A)^*$ is generated by separable sub σ -algebras of the algebras $\sigma(B)$ where $B \in A^*$. In particular we can choose separable σ -algebras $d'(A, B) \supset c(B)$ in $\sigma(B)$ for every $B \in A^*$ such that $\langle d'(A, B)|B \in A^* \rangle \supseteq d(A)^*$. For a set $B \subseteq [k]$ with $|B| = t - 1$ we define $d'(B)$ as the σ -algebra generated by all the algebras in the form of $d'(C, D)^\pi$, where $\pi \in S_k$, $D^\pi = B$, $|C| = |D| + 1$ and $D \subseteq C$. Since $d'(C, D)^\pi \subseteq \sigma(D)^\pi = \sigma(B)$ we have that $d'(B) \subseteq \sigma(B)$. Furthermore we have that $d'(B)^\pi = d'(B^\pi)$ for every $\pi \in S_k$.

Now let $d(A) := \langle d'(B) \mid B \subseteq A \rangle$. the second requirement in the lemma is trivial by definition. We prove the first one. The elements of $d(A)$ can be approximated by finite unions of intersections of the form $\bigcap_{B \subseteq A} T_B$ where $T_B \in d'(B)$ and so it is enough to prove the statement if R is such an intersection. Let $Q = \bigcap_{B \subseteq A, B \neq A} T_B$. Now

$$E(R|\langle d(B)|B \in A^* \rangle) = E(R|\langle d'(B)|B \subseteq A, B \neq A \rangle).$$

By the basic property of the conditional expectation (see Appendix) :

$$\begin{aligned} E(R|\langle d'(B)|B \subseteq A, B \neq A \rangle) &= E(T_A|\langle d'(B)|B \subseteq A, B \neq A \rangle)\chi_Q = E(T_A|\sigma(A)^*)\chi_Q = \\ &= E(R|\sigma(A)^*). \end{aligned}$$

■

Proof of Theorem 6 We construct the algebras $l(A)$ in the following steps. For each non-empty subset $A \subseteq [k]$ we choose an atomless separable σ -algebra $c(A) \subseteq \sigma(A)$ containing a $\sigma(A)^*$ -random r -partition for every r . We also assume that $\mathbf{H} \in c([k])$. Applying Lemma 2.12 for the previous system of separable σ -algebras $c(A)$ we obtain the σ -algebras $d(A)$. By Lemma 2.11 and the permutation invariance property of the previous lemma, $S_{[r]}$ acts freely on $d([r])^*$. Hence using Lemma 2.9, for every $\emptyset \neq A \in [k]$ we can choose an independent complement $l([r])$ for $d([r])^* = \langle d(B)|B \in [r]^* \rangle$ in $d([r])$ such that $l([r])$ is element-wise invariant under the action of $S_{[r]}$. The algebras $l([r])$ are independent from $\sigma([r])^*$ since $\mu(R) = E(R|d([r])^*) = E(R|\sigma([r])^*)$ for every $R \in l([r])$. Now we define $l(A)$, where $|A| = r$ by $l(A) = l([r])^\pi$ for some $\pi \in S_k$, $\pi([r]) = A$. Note that $l(A)$ does not depend on the choice of π . By Lemma 4.1 of the Appendix we have maps $F_{[r]} : X \rightarrow [0, 1]$ such that F^{-1} defines a measure algebra isomorphism between $\mathcal{M}([0, 1], \mathcal{B}, \lambda)$ and $\mathcal{M}(\mathbf{X}, l[r], \mu)$. Let $F_A = \pi^{-1} \circ F_{[r]}$, where π maps $[r]$ to A . Again, $F_{[r]}$ does not depend on the particular choice of the permutation π . ■

Now let $\mathcal{S} = \{l(A), F_A\}_{\emptyset \neq A \subseteq [k]}$ be a separable realization of $\mathbf{X}$ and $k < n$ be a natural number. Let $B \subset [n], |B| = r \leq k$ and $\pi \in S_n$ be a permutation that maps $[r]$ to B . Let $l(B) \subseteq \mathbf{X}^n$ be defined as $l([r])^\pi$. If we choose a $\pi' \in S_n$ that also maps $[r]$ into B then $(\pi)^{-1} \circ \pi'$ permutes $[r]$ hence fixes $l([r])$. Therefore $l(B)$ does not depend on the choice of π . Let F_B defined as $\pi^{-1} \circ F_{[r]}$. We have the following lemma.

Lemma 2.13 *The system $\tilde{\mathcal{S}} = \langle l(B), F_B \rangle_{\emptyset \neq B \subseteq [n], |B| \leq k}$ is a separable realization of degree k on X^n . If $B \subset [n], |B| = r \leq k$ and $f : [r] \rightarrow B$ is a bijection then let $p_B : \mathbf{X}^n \rightarrow \mathbf{X}^B$ the natural projection and $p_f : \mathbf{X}^{[r]} \rightarrow \mathbf{X}^B$ is the natural isomorphism. Then $l(B) = p_B^{-1}(p_f(l([r])))$.*

By Lemma 2.4 and Lemma 4.2 we have the following lifting lemma as well.

Lemma 2.14 *The map $F : X^k \rightarrow [0, 1]^{2^k - 1}$, $F = \oplus_{\emptyset \neq A \subseteq [k]} F_A$ defines an isomorphism between the measure algebra of $\mathcal{M}(\mathbf{X}^k, \langle l(A) \mid \emptyset \neq A \subseteq [k] \rangle, \mu)$ and the Lebesgue measure algebra $\mathcal{M}([0, 1]^{2^k - 1}, \mathcal{B}, \lambda)$. Similarly, $\tilde{F} : \mathbf{X}^n \rightarrow [0, 1]^{\sum_{i=1}^k \binom{n}{i}}$, $\tilde{F} = \oplus_{\emptyset \neq B \subseteq [n], |B| \leq k} F_B$ defines an isomorphism between the measure algebra $\mathcal{M}(\mathbf{X}^n, \langle l(B) \mid \emptyset \neq B \subseteq [n], |B| \leq k \rangle, \mu)$ and the Lebesgue measure algebra $\mathcal{M}([0, 1]^{\sum_{i=1}^k \binom{n}{i}}, \mathcal{B}, \lambda)$.*

3 Applications for Hypergraphs

3.1 Hypergraph homomorphisms and convergence

Recall that a k -uniform **hypergraph** H is a system of k element subsets (**edges**) denoted by $E(H)$ of a set V (**node set**). A k -uniform hypergraph can be represented as a subset $S_H \subset V^k$ such that $(x_1, x_2, \dots, x_k) \in S$ if and only if $\{x_1, x_2, \dots, x_k\} \in E(H)$. Note that S_H is invariant under the action of S_k on V^k . For any hypergraph we have an underlying $(k-1)$ -dimensional simplicial complex $\Sigma(H)$ consisting of the subsets of the k -edges.

Suppose that K is a finite k -uniform hypergraph on the node set $[n] := \{1, 2, \dots, n\}$ and H is a k -uniform hypergraph on the node set V . Then a map $f : [n] \rightarrow V$ is a **homomorphism** if f maps edges to edges. If H is finite then $\text{hom}(K, H)$ is the number of homomorphism from K to H . Denote by $t(K, H)$ the probability that a random map $g : [n] \rightarrow V$ is a (K, H) -homomorphism, that is

$$t(K, H) = \frac{\text{hom}(K, H)}{|V|^n}.$$

If H is not necessarily finite then $T(K, H) \subset V^n$ denotes the (K, H) -**homomorphism set**, where $(x_1, x_2, \dots, x_n) \in T(K, H)$ if $1 \rightarrow x_1, 2 \rightarrow x_2, \dots, n \rightarrow x_n$ defines a homomorphism. Clearly $|T(K, H)| = \text{hom}(K, H)$. Note that

$$T(K, H) = \bigcap_{E \in E(K)} p_E^{-1}(p_f(S_H)),$$

where $f : [k] \rightarrow B$ is a bijection (see Lemma 2.14).

Definition 3.1 *We say that a sequence of k -uniform hypergraphs $\{H_i\}_{i=1}^\infty$ is **convergent** if for every fixed finite k -uniform hypergraph K $\lim_{i \rightarrow \infty} t(K, H_i)$ exists.*

Let $\{X_i\}_{i=1}^\infty$ be finite sets and $H_i \subset X_i^k$ be k -uniform directed hypergraphs, that is a sequence of S_k -invariant sets $S_{H_i} \subset X_i^k$ is given. As in the Section 2.1, let X be the ultralimit of the sets X_i . Then $\mathbf{H} := [\{S_{H_i}\}_{i=1}^\infty] \subset \mathcal{P}(X^k)$ is the **ultralimit hypergraph**, an S_k -invariant set corresponding to an actual hypergraph on the node set X . We can define its homomorphism set as

$$T(K, \mathbf{H}) := \bigcap_{E \in E(K)} p_E^{-1}(p_f(\mathbf{H})) .$$

Then

$$T(K, \mathbf{H}) = [\{T(K, H_i)\}_{i=1}^\infty] \subset \mathcal{P}(\mathbf{X}^n) .$$

Clearly, $\mu^n(T(K, \mathbf{H})) = \lim_\omega t(K, H_i)$, where μ^n denotes the ultralimit measure on $\mathbf{X}^n$. Thus if $\{H_i\}_{i=1}^\infty$ is a convergent sequence of hypergraphs then:

$$\mu^n(T(K, \mathbf{H})) = \lim_{i \rightarrow \infty} t(K, H_i) .$$

3.2 The Hypergraph Removal Lemma

Lemma 3.1 (Infinite Removal Lemma) *Let $\mathbf{H}$ be an S_k -invariant measurable subset of $\mathbf{X}^k$. Then there exists an S_k -invariant nullset $\mathbf{I} \subseteq \mathbf{H}$ such that for every k -uniform hypergraph K either $T(K, \mathbf{H} \setminus \mathbf{I}) = \emptyset$ or $|T(K, \mathbf{H} \setminus \mathbf{I})| > 0$.*

Proof. Let us consider the separable realization $\mathcal{S}$ of $\mathbf{H}$ and the corresponding measurable equivalence $F : \mathbf{X} \rightarrow [0, 1]^{2^k-1}$. For some Lebesgue measurable set $Q \subseteq [0, 1]^{2^k-1}$ we have that $|F^{-1}(Q) \Delta \mathbf{H}| = 0$. Since

$$F^{-1}(Q^\pi) \Delta \mathbf{H}^\pi = (F^{-1}(Q) \Delta \mathbf{H})^\pi$$

we may suppose that Q is S_k -invariant. By Lebesgue's Density Theorem, almost all points of Q are density points. Let D denote the (S_k -invariant) set of density points in Q and let $S := F^{-1}(D)$. Notice that the group S_k acts on $[0, 1]^{2^k-1}$ the following way. Let $A_1, A_2, \dots, A_{2^k-1}$ be a list of non-empty subsets of $[k]$. Then

$$(y_{A_1}, y_{A_2}, \dots, y_{A_{2^k-1}})^\pi = (y_{\pi^{-1}(A_1)}, y_{\pi^{-1}(A_2)}, \dots, y_{\pi^{-1}(A_{2^k-1})}) .$$

By the invariance property of the separable realization, the maps F_A commutes with the S_k -action that is $\pi \circ F_A = F_A \circ \pi$. Also, let $B_1, B_2, \dots, B_r$, ($r = \sum_{i=1}^k \binom{n}{i}$) be the list of

non-empty subsets of $[n]$ of size at most k , then S_n acts on $[0, 1]^r$ by

$$(y_{B_1}, y_{B_2}, \dots, y_{B_r})^\pi = (y_{\pi^{-1}(B_1)}, y_{\pi^{-1}(B_2)}, \dots, y_{\pi^{-1}(B_r)}).$$

Again, by the invariance property of the lifting $\rho \circ F_B = F_B \circ \rho$, for any $B \subset [n]$, $|B| \leq k$, $\rho \in S_n$. For $B \subset [n]$, $|B| = k$ a bijection $f : [k] \rightarrow B$ induces a measurable isomorphism $L_f : [0, 1]^{2^k-1} \rightarrow [0, 1]^{r(B)}$, where $r(B)$ denotes the set of non-empty subsets of B . Let $L_B : [0, 1]^r \rightarrow [0, 1]^{r(B)}$ be the natural projection. Then by the invariance property of the lifting

$$p_B^{-1}(p_f(\mathbf{X}^k)) = \tilde{F}^{-1}(L_B^{-1}(L_f([0, 1]^{2^k-1}))). \quad (4)$$

That is for any k -regular hypergraph K

$$T(K, \mathbf{S}) = \cap_{E \in E(K)} p_E^{-1}(p_f(\mathbf{S})) = \tilde{F}^{-1}(\cap_{E \in E(K)} L_E^{-1}(L_f(D))).$$

Since each point of D is a density point, each point of $L_E^{-1}(L_f(D))$ is a density point for any $E \in E(K)$. Thus $\cap_{E \in E(K)} L_E^{-1}(L_f(D))$ is either empty or is of positive measure. Consequently, $T(K, \mathbf{S})$ is either empty or is of positive measure as well. Choosing $\mathbf{I} = H \setminus \mathbf{S}$, we obtain that $T(K, H \setminus \mathbf{I}) = T(K, H \cap \mathbf{S})$ is either empty or is of positive measure (note that $\mu(T(K, H \cap \mathbf{S})) = \mu(T(K, \mathbf{S}))$ and $T(K, H \cap \mathbf{S}) \subseteq T(K, \mathbf{S})$). ■

Theorem 7 (Hypergraph Removal Lemma) *For every k -uniform hypergraph K and constant $\epsilon > 0$ there exists a number $\delta = \delta(K, \epsilon)$ such that for any k -uniform hypergraph H on the node set X with $t(K, H) < \delta$ there is a subset L of $E(H)$ with $|L| \leq \epsilon \binom{|X|}{k}$ such that $t(K, H \setminus L) = 0$. ([2], [4], [8], [11])*

Proof. We proceed by contradiction. Let K be a fixed hypergraph and $\epsilon > 0$ be a fixed number for which the theorem fails. This means that there is a sequence of hypergraphs H_i on the sets X_i such that $\lim_{i \rightarrow \infty} t(K, H_i) = 0$ but in each H_i there is no set L with the required property. Let us represent the hypergraphs by symmetric subsets S_{H_i} of X_i^k and again let $\mathbf{H} \subseteq \mathbf{X}^k$ denote the ultralimit of them. Then $\mu(T(K, \mathbf{H})) = \lim_\omega t(K, H_i) = 0$ and thus by the previous lemma there is a zero measure S_k -invariant set $\mathbf{I} \subseteq \mathbf{X}^k$ such that $T(K, \mathbf{H} \setminus \mathbf{I}) = \emptyset$. By the definition of nullsets, for any $\epsilon_1 > 0$ there exists an ultralimit set $\mathbf{J} \subset \mathbf{X}^k$ such that $\mathbf{I} \subset \mathbf{J}$ and $\mu(\mathbf{J}) \leq \epsilon_1$. We can suppose that $\mathbf{J}$ is S_k -invariant as well. Let $[\{J_i\}_{i=1}^\infty] = \mathbf{J}$, then for ω -almost all i , J_i is S_k -invariant, $|J_i| \leq \epsilon_1 |X_i|^k$ and $T(K, H_i \setminus L_i) = \emptyset$, where L_i is the set of edges $\{x_1, x_2, \dots, x_k\}$ such that $(x_1, x_2, \dots, x_k) \in J_i$. Clearly, $|L_i| \leq |J_i|$, hence if ϵ_1 is small enough then $|L_i| \leq \epsilon \binom{|X_i|}{k}$ leading to a contradiction. ■

3.3 The Hypergraph Limit Object

In this section we introduce the notion of **hypergraphons** (see [6] and [1] for **graphons**). Let $W : [0, 1]^{2^k-1} \rightarrow \{0, 1\}$ be a Lebesgue measurable function. We call such functions **directed hypergraphons**. As in the previous subsection we consider the S_k -action on $[0, 1]^{2^k-1}$ and call the S_k -invariant directed hypergraphons just hypergraphons. Now we introduce the homomorphism density of a hypergraph into a hypergraphon. Let K be a k -uniform hypergraph and $W : [0, 1]^{2^k-1} \rightarrow \mathbb{R}$ be a hypergraphon. Let $C_K = \{C_1, C_2, \dots, C_s\}$ be the set of non-empty elements of the simplicial complex of K .

Example: If $K = \{\{1, 2, 3\}, \{2, 3, 4\}\}$ then

$$C_K = \{\{1\}, \{2\}, \{3\}, \{4\}, \{12\}, \{13\}, \{23\}, \{24\}, \{34\}, \{1, 2, 3\}, \{2, 3, 4\}\}.$$

For each edge $E \in E(K)$ we fix a bijection $s_E : [k] \rightarrow E$. Then the **homomorphism density** of K in W is defined as

$$t(K, W) := \int_0^1 \int_0^1 \dots \int_0^1 \prod_{E \in E(K)} W_{\mathbf{H}}(x_{s_E(A_1)}, x_{s_E(A_2)}, \dots, x_{s_E(A_{2^k-1})}) dx_{C_1} dx_{C_2} \dots dx_{C_s}. \quad (5)$$

Now let $\{X_i\}_{i=1}^\infty$ be finite sets and H_i be k -uniform directed hypergraphs on X_i . Let $\mathbf{H} \subset \mathbf{X}^k$ be their ultralimit hypergraph. Let $F : X^k \rightarrow [0, 1]^{2^k-1}$ be separable realization and $Q \subset [0, 1]^{2^k-1}$ be a S_k -invariant measurable set such that $\mu(F^{-1}(Q) \triangle \mathbf{H}) = 0$. Then we define $W_{\mathbf{H}}$ as the characteristic function of Q . Clearly, $W_{\mathbf{H}}$ is an hypergraphon. Now we can state our main theorem.

Theorem 8 (Main Theorem) *Let $\{H_i\}_{i=1}^\infty$ be a sequence of k -uniform hypergraphs as above and let K be a fixed k -uniform hypergraph on the vertex set $[n]$. Then*

$$\lim_{\omega} t(K, H_i) = t(K, W_{\mathbf{H}}).$$

Proof. Applying the Equation (4) we obtain that

$$\mu(T(K, H)) = \text{Vol} \left(\cap_{E \in E(K)} L_E^{-1}(L_f(Q)) \right).$$

Hence

$$\mu(T(K, H)) = \int_0^1 \int_0^1 \dots \int_0^1 \prod_{E \in E(K)} \Psi_E dx_{B_1} dx_{B_2} \dots dx_{B_r},$$

where Ψ_E is the characteristic function of $L_E^{-1}(L_f(Q))$. Clearly,

$$\Psi_E(x_{B_1}, x_{B_2}, \dots, x_{B_r}) = W_{\mathbf{H}}(x_{s_E(A_1)}, x_{s_E(A_2)}, \dots, x_{s_E(A_{2^k-1})}).$$

Since $\prod_{E \in E(K)} \Psi_E$ depends only on the variables associated to the elements of the simplicial complex of K , the Theorem follows. $\blacksquare$

The following theorem is an immediate corollary of the previous one.

Theorem 9 *If $\{H_i\}_{i=1}^\infty$ is a convergent sequence of k -uniform hypergraphs then there exists a $2^k - 1$ variable hypergraphon W such that $\lim_{i \rightarrow \infty} t(K, H_i) = t(K, W)$ for every k -uniform hypergraph K .*

Remark: One can introduce the notion of a **projected hypergraphon** $\widetilde{W}_{\mathbf{H}}$ which is the projection of a hypergraphon to the first $2^k - 2$ coordinates, where the last coordinate is associated to $[k]$ itself. That is

$$\widetilde{W}_{\mathbf{H}}(x_{A_1}, x_{A_2}, \dots, x_{A_{2^k-2}}) = \int_0^1 W_{\mathbf{H}}(x_{A_1}, x_{A_2}, \dots, x_{A_{2^k-1}}) dx_{A_{2^k-1}}.$$

That is $\widetilde{W}_{\mathbf{H}}$ is a $[0, 1]$ -valued function which is symmetric under the induced S_k -action of its coordinates. By the classical Fubini-theorem we obtain that using the notation of the previous theorem:

$$\begin{aligned} \lim_{i \rightarrow \infty} t(K, H_i) &= \\ &= \int_0^1 \int_0^1 \dots \int_0^1 \prod_{E \in E(K)} \widetilde{W}_{\mathbf{H}}(x_{s_E(A_1)}, x_{s_E(A_2)}, \dots, x_{s_E(A_{2^k-2})}) dx_{C_1} dx_{C_2} \dots dx_{C_t}, \end{aligned}$$

where $C_1, C_2, \dots, C_t$ is the list of the at most $k - 1$ -dimensional simplices in K . Note that in the case $k = 2$ it is just the graph limit formula of [6].

3.4 The Hypergraph Regularity Lemma

First we need some definitions. Let X be a finite set, then $K_r(X)$ denotes the complete r -uniform hypergraph on X . An l -**hyperpartition** $\mathcal{H}$ is a family of partition $K_r(X) = \cup_{j=1}^l P_r^j$, where P_r^j is an r -uniform hypergraph, for $1 \leq r \leq k$. We call $\mathcal{H}$ **δ -equitable** if for any $1 \leq r \leq k$ and $1 \leq i < j \leq l$:

$$\frac{||P_r^i| - |P_r^j||}{|K_r(X)|} < \delta.$$

An l -hyperpartition $\mathcal{H}$ induces a partition on $K_k(X)$ the following way.

- Two elements $\underline{a}, \underline{b} \in K_k(X)$, $\underline{a} = \{a_1, a_2, \dots, a_k\}$, $\underline{b} = \{b_1, b_2, \dots, b_k\}$ are equivalent if there exists a permutation $\sigma \in S_k$ such that for any subset $A = \{i_1 < i_2 < \dots < i_{|A|}\} \in [k]$, $\{a_{i_1}, a_{i_2}, \dots, a_{i_{|A|}}\}$ and $\{b_{\sigma(i_1)}, b_{\sigma(i_2)}, \dots, b_{\sigma(i_{|A|})}\}$ are both in the same $P_{|A|}^j$ for some $1 \leq j \leq l$.

It is easy to see that this defines an equivalence relation and thus it results in a partition $\cup_{j=1}^t C_j$ of $K_k(X)$ into $\mathcal{H}$ -cells. A **cylinder intersection** $L \subset K_r(X)$ is an r -uniform hypergraph defined the following way. Let $B_1, B_2, \dots, B_r$ be $r - 1$ uniform hypergraphs on X , then an r -edge $\{a_1, a_2, \dots, a_r\}$ is in L if there exists a permutation $\tau \in S_r$ such that

$$\{a_{\sigma(1)}, a_{\sigma(2)}, \dots, a_{\sigma(i-1)}, a_{\sigma(i+1)}, \dots, a_{\sigma(r)}\} \in B_i.$$

As in the graph case, we call an r -uniform hypergraph G ϵ -regular if

$$\left| \frac{|G|}{|K_r(X)|} - \frac{|G \cap L|}{|L|} \right| \leq \epsilon,$$

for each cylinder intersection L , where $|L| \geq \epsilon |K_r(X)|$. Now we are ready to state the hypergraph regularity lemma for k -uniform hypergraphs (see [2], [4], [9], [11]).

Theorem 10 (Hypergraph regularity lemma) *Let fix a constant $k > 0$. Then for any $\epsilon > 0$ and $F : \mathbb{N} \rightarrow (0, 1)$ there exists constants $c = c(\epsilon, F)$ and $N_0(\epsilon, F)$ such that if H is a k -uniform hypergraph on a set X , $|X| \geq N_0(\epsilon, F)$, then there exists an $F(l)$ -equitable l -hyperpartition $\mathcal{H}$ for some $1 < l \leq c$ such that*

- Each P_j^r is $F(l)$ -regular.
- $|H \Delta T| \leq \epsilon \binom{|X|}{k}$ where T is the union of some $\mathcal{H}$ -cells.

Proof. Suppose that the Theorem does not hold for some $\epsilon > 0$ and $F : \mathbb{N} \rightarrow (0, 1)$. That is there exists a sequence of k -uniform hypergraphs H_i without having $F(j)$ -equitable j -hyperpartitions for any $1 < j \leq i$ satisfying the conditions of our Theorem. Let us consider their ultralimit $[\{S_{H_i}\}_{i=1}^\infty] = \mathbf{H} \subset \mathbf{X}^k$. Similarly to the proof of the Removal Lemma we formulate an infinite version of the Regularity Lemma as well.

Let $K_r(\mathbf{X})$ denote the complete r -uniform hypergraph on X , that is the set of points $(x_1, x_2, \dots, x_r) \in \mathbf{X}^r$ such that $x_i \neq x_j$ if $i \neq j$. Clearly $K_r(\mathbf{X}) \subset \mathbf{X}^r$ is measurable and $\mu_{[r]}(K_r(\mathbf{X})) = 1$. An r -uniform hypergraph on $\mathbf{X}$ is an S_r -invariant measurable subset of $K_r(\mathbf{X})$. An l -hyperpartition $\tilde{\mathcal{H}}$ is a family of partitions $K_r(\mathbf{X}) = \cup_{j=1}^l \mathbf{P}_r^j$, where $\mathbf{P}_r^j$ is an r -uniform hypergraph for $1 \leq r \leq k$. Again, an l -hyperpartition induces a partition of $K_k(\mathbf{X})$ into $\tilde{\mathcal{H}}$ -cells exactly the same way as in the finite case. It is easy to see that each $\tilde{\mathcal{H}}$ -cell is measurable.

Proposition 3.1 (Hypergraph Regularity Lemma, infinite version) *For any $\epsilon > 0$, there exists a 0-equitable l -hyperpartition (where l depends on $\mathbf{H}$) $\tilde{\mathcal{H}}$ such that*

- Each $\mathbf{P}_r^j$ is in $\sigma([r])^*$.
- $\mu_{[k]}(H \Delta T) \leq \epsilon$, where T is a union of some $\tilde{\mathcal{H}}$ -cells.

Proof. Let $\mathcal{S}$ be a separable realization for $\mathbf{H}$ and $Q \subseteq [0, 1]^{2^k-1}$ be an S_k -invariant subset such that $\mu_{[k]}(F^{-1}(Q) \Delta \mathbf{H}) = 0$. Since Q is a Lebesgue-measurable set, there exists some $l > 0$ such that $\text{Vol}_{2^k-1}(Q \Delta Z) < \epsilon$, where Z is a union of l -boxes. Recall that an l -box is a product set in the form

$$\left(\frac{i_1}{l}, \frac{i_1+1}{l} \right) \times \left(\frac{i_2}{l}, \frac{i_2+1}{l} \right) \times \dots \times \left(\frac{i_{2^k-1}}{l}, \frac{i_{2^k-1}+1}{l} \right).$$

By the usual symmetrization argument we may suppose that the set Z is invariant under the S_k -action on the l -boxes. Since the measure of points $(x_1, x_2, \dots, x_{2^k}) \in [0, 1]^{2^k-1}$ such that $x_s = x_t$ for some $s \neq t$ is zero, we may also suppose that in each box in Z , $i_s \neq i_t$ if $s \neq t$. Let $Z = \cup_{m=1}^q O_m$, where O_m is an S_k -orbit of boxes. That is $O_m = \cup_{\pi \in S_k} \pi(D)$ for some l -box D . By the previous condition $\pi_1(D) \neq \pi_2(D)$, if $\pi_1 \neq \pi_2$, hence each O_m is the disjoint union of exactly $k!$ l -boxes. Then $\mu_{[k]}(F^{-1}(Q) \triangle F^{-1}(Z)) < \epsilon$, where $F^{-1}(Z) = \cup_{m=1}^q F^{-1}(O_m)$. For each $1 \leq r \leq k$ we consider the partition $\mathbf{X}^r = \cup_{j=1}^l \mathbf{P}_r^j$, where $\mathbf{P}_r^j = F_{[r]}^{-1}(\frac{j-1}{l}, \frac{j}{l})$. We call the resulting l -hyperpartition $\tilde{\mathcal{H}}$. Note that by the S_r -invariance of the separable realization each $\mathbf{P}_r^j$ is an r -uniform hypergraph and also $\mathbf{P}_r^j \in \sigma([r])^*$.

Lemma 3.2 $\mathbf{C}$ is an $\tilde{\mathcal{H}}$ -cell if and only if $\mathbf{C} = F^{-1}(\cup_{\pi \in S_k} \pi(D))$, where D is an l -box in $[0, 1]^{2^k-1}$.

Proof. By definition $(a_1, a_2, \dots, a_k) \in \mathbf{X}^k$ and $(b_1, b_2, \dots, b_k) \in \mathbf{X}^k$ are in the same $\tilde{\mathcal{H}}$ -cell if and only if there exists $\pi \in S_k$ such that for any $A \subseteq [k]$ $(a_{i_1}, a_{i_2}, \dots, a_{i_{|A|}})$ and $(b_{i_{\pi(1)}}, b_{i_{\pi(2)}} \dots, b_{i_{\pi(|A|)}})$ are in the same $\mathbf{P}_r^j$. That is $F_A(a_1, a_2, \dots, a_k)$ and $F_A(b_{\pi(1)}, b_{\pi(2)}, \dots, b_{\pi(k)})$ are in the same l -box. ■

Since $\mu_{[k]}(H \triangle \cup_{j=1}^q F^{-1}(O_m)) < \epsilon$, our Proposition follows. ■

Now we return to the proof of the Hypergraph Regularity Lemma. First pick an r -hypergraph $\tilde{\mathbf{P}}_r^j$ on $\mathbf{X}$ such that $\mu_{[r]}(\tilde{\mathbf{P}}_r^j \triangle \mathbf{P}_r^j) = 0$, $\tilde{\mathbf{P}}_r^j \in \mathcal{P}_{[r]}$ and $\cup_{j=1}^l \tilde{\mathbf{P}}_r^j = K_r(\mathbf{X})$. Let $[\{S_{P_{r,i}^j}\}_{i=1}^\infty] = \tilde{\mathbf{P}}_r^j$. Then for ω -almost all indices $\cup_{j=1}^l P_{r,i}^j = K_r(X_i)$ is an $F(l)$ -equitable l -partition and $|H_i \triangle \cup_{m=1}^q C_m^i| < \epsilon$ for the induced $\mathcal{H}$ -cell approximation. Here $\cup_{m=1}^q \tilde{\mathbf{C}}_m$ is the $\tilde{\mathcal{H}}$ -cell approximation with respect to the l -hyperpartitions $\cup_{j=1}^l \tilde{\mathbf{P}}_r^j = K_r(\mathbf{X})$ and $[\{S_{C_m^i}\}_{i=1}^\infty] = \tilde{\mathbf{C}}_m$.

The only thing remained to be proved is that for ω -almost all indices i the resulting l -hyperpartitions are $F(l)$ -regular. If it does not hold then there exists $1 \leq r \leq k$ and $1 \leq j \leq l$ such that for almost all i there exists a cylinder intersection $W_i \subset K_r(X_i)$, $|W_i| \geq \epsilon |X_i|$, such that

$$\left| \frac{|P_{r,i}^j|}{|K_r(X_i)|} - \frac{|P_{r,i}^j \cap W_i|}{|W_i|} \right| > \epsilon. \quad (6)$$

Let $\mathbf{W} = [\{S_{W_i}\}_{i=1}^\infty]$. Then $W \subset \cup_{B \subsetneq [r]} \sigma(B)$. Hence $\tilde{\mathbf{P}}_r^j$ and $\mathbf{W}$ are independent sets. However, by (6)

$$\mu_{[r]}(\tilde{\mathbf{P}}_r^j) \mu_{[r]}(\mathbf{W}) \neq \mu_{[r]}(\tilde{\mathbf{P}}_r^j \cup \mathbf{W}),$$

leading to a contradiction. ■

4 Appendix on basic measure theory

In this section we collect some of the basic results of measure theory we frequently use in our paper.

Separable measure spaces: Let $(X, \mathcal{A}, \mu)$ be a probability measure space. Then we call $A, A' \in \mathcal{A}$ equivalent if $\mu(A \Delta A') = 0$. The equivalence classes form a complete metric space, where $d([A], [B]) = \mu(A \Delta B)$. These classes form a Boolean-algebra as well, called the **measure algebra** $\mathcal{M}(X, \mathcal{A}, \mu)$. We say that $(X, \mathcal{A}, \mu)$ is a **separable** measure space if $\mathcal{M}(X, \mathcal{A}, \mu)$ is a separable metric space. It is important to note that if $(X, \mathcal{A}, \mu)$ is separable and atomless, then its measure algebra is isomorphic to the measure algebra of the standard Lebesgue space $([0, 1], \mathcal{B}, \lambda)$, where $\mathcal{B}$ is the σ -algebra of Borel sets (see e.g. [3]). We use the following folklore version of this theorem.

Lemma 4.1 *If $(X, \mathcal{A}, \mu)$ is a separable and atomless measure algebra, then there exists a map $f : X \rightarrow [0, 1]$ such that $f^{-1}(\mathcal{B}) \subset \mathcal{A}$, $\mu(f^{-1}(U)) = \lambda(U)$ for any $U \in \mathcal{B}$ and for any $L \in \mathcal{A}$ there exists $M \in \mathcal{B}$ such that L is equivalent to $f^{-1}(M)$.*

Proof. Let I_0 denote the interval $[0, \frac{1}{2}]$, $I_1 = [\frac{1}{2}, 1]$. Then let $I_{0,0} = [0, \frac{1}{4}]$, $I_{0,1} = [\frac{1}{4}, \frac{1}{2}]$, $I_{1,0} = [\frac{1}{2}, \frac{3}{4}]$, $I_{1,1} = [\frac{3}{4}, 1]$. Recursively, we define the dyadic intervals $I_{\alpha_1, \alpha_2, \dots, \alpha_k}$, where $(\alpha_1, \alpha_2, \dots, \alpha_k)$ is a 0-1-string. Let T be the Boolean-algebra isomorphism between the measure algebra of $(X, \mathcal{A}, \mu)$ and the measure algebra of $([0, 1], \mathcal{B}, \lambda)$. Then we have disjoint sets $U_0, U_1 \in \mathcal{A}$ such that $T([U_0]) = [I_0]$, $T([U_1]) = [I_1]$. Clearly $\mu(X \setminus (U_0 \cup U_1)) = 0$. Similarly, we have disjoint subsets of U_0 , $U_{0,0}$ and $U_{0,1}$ such that $T([U_{0,0}]) = [I_{0,0}]$ and $T([U_{0,1}]) = [I_{0,1}]$. Recursively, we define $U_{\alpha_1, \alpha_2, \dots, \alpha_k} \in \mathcal{A}$ such that $U_{\alpha_1, \alpha_2, \dots, \alpha_{k-1}, 0}$ and $U_{\alpha_1, \alpha_2, \dots, \alpha_{k-1}, 1}$ are disjoint and $T([U_{\alpha_1, \alpha_2, \dots, \alpha_k}]) = I_{\alpha_1, \alpha_2, \dots, \alpha_k}$. The set of points in X which are not included in some $U_{\alpha_1, \alpha_2, \dots, \alpha_k}$ for some $k > 0$ has measure zero. Now define

$$f(p) := \bigcap_{k=1}^{\infty} I_{\alpha_1, \alpha_2, \dots, \alpha_k},$$

where for each $k \geq 1$, $p \in U_{\alpha_1, \alpha_2, \dots, \alpha_k}$. It is easy to see that f satisfies the conditions of our lemma. ■

Generated σ -algebras: Let $(X, \mathcal{C}, \mu)$ be a probability measure space and $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k$ be sub- σ -algebras. Then we denote by $\langle \mathcal{A}_i \mid 1 \leq i \leq k \rangle$ the generated σ -algebra that is the smallest sub- σ -algebra of $\mathcal{C}$ containing the $\mathcal{A}_i$'s. Then the equivalence classes

$$[\bigcup_{j=1}^n (A_1^j \cap A_2^j \cap \dots \cap A_k^j)],$$

where $A_i^j \in \mathcal{A}_i$ and $(A_1^s \cap A_2^s \cap \dots \cap A_k^s) \cap (A_1^t \cap A_2^t \cap \dots \cap A_k^t) = \emptyset$ if $s \neq t$ form a dense subset in the measure algebra $\mathcal{M}(X, \langle \mathcal{A}_i \mid 1 \leq i \leq k \rangle, \mu)$ with respect to the metric defined above (see [3]).

Independent subalgebras and product measures: The sub- σ -algebras $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k \subset \mathcal{C}$ are **independent** subalgebras if

$$\mu(A_1)\mu(A_2)\dots\mu(A_k) = \mu(A_1 \cap A_2 \cap \dots \cap A_k),$$

if $A_i \in \mathcal{A}_i$.

Lemma 4.2 *Let $\mathcal{A}_1, \mathcal{A}_2, \dots, \mathcal{A}_k \subset \mathcal{C}$ be independent subalgebras as above and $f_i : X \rightarrow [0, 1]$ be maps such that f_i^{-1} defines isomorphisms between the measure algebras $\mathcal{M}(X, \mathcal{A}_i, \mu)$ and $\mathcal{M}([0, 1], \mathcal{B}, \lambda)$. Then the map $F^{-1}, F = \bigoplus_{i=1}^k f_i : X \rightarrow [0, 1]^k$ defines an isomorphism between the measure algebras $\mathcal{M}(X, \langle \mathcal{A}_i \mid 1 \leq i \leq k \rangle, \mu)$ and $\mathcal{M}([0, 1]^k, \mathcal{B}^k, \lambda^k)$.*

Proof. Observed that

$$\mu(F^{-1}(\cup_{i=1}^s [A_1^i \times \dots \times A_k^i])) = \sum_{i=1}^s \lambda^k[A_1^i \times \dots \times A_k^i]$$

whenever $\{A_1^i \times \dots \times A_k^i\}_{i=1}^s$ are disjoint product sets. Hence F^{-1} defines an isometry between dense subsets of the two measure algebras. ■

Radon-Nykodym Theorem: Let $(X, \mathcal{A}, \mu)$ be a probability measure space and ν be an absolutely continuous measure with respect to μ . That is if $\mu(A) = 0$ then $\nu(A) = 0$ as well. Then there exists an integrable $\mathcal{A}$ -measurable function f such that

$$\nu(A) = \int_A f d\mu$$

for any $A \in \mathcal{A}$.

Conditional expectation: Let $(X, \mathcal{A}, \mu)$ be a probability measure space and $\mathcal{B} \subset \mathcal{A}$ be a sub- σ -algebra. Then by the Radon-Nykodym-theorem for any integrable $\mathcal{A}$ -measurable function f there exists an integrable $\mathcal{B}$ -measurable function $E(f \mid \mathcal{B})$ such that

$$\int_B E(f \mid \mathcal{B}) d\mu = \int_B f d\mu,$$

if $B \in \mathcal{B}$. The function $E(f \mid \mathcal{B})$ is called the conditional expectation of f with respect to $\mathcal{B}$. It is unique up to a zero-measure perturbation. Note that if $a \leq f(x) \leq b$ for almost all $x \in X$, then $a \leq E(f \mid \mathcal{B})(x) \leq b$ for almost all $x \in X$ as well. Also, if g is a bounded $\mathcal{B}$ -measurable function, then

$$E(fg \mid \mathcal{B}) = E(f \mid \mathcal{B})g \quad \text{almost everywhere.}$$

The map $f \rightarrow E(f, \mathcal{B})$ extends to a Hilbert-space projection $E : L^2(X, \mathcal{A}, \mu) \rightarrow L^2(X, \mathcal{B}, \mu)$.

Lebesgue density theorem: Let $A \in \mathbb{R}^n$ be a measurable set. Then almost all points $x \in A$ is a **density point**. The point x is a density point if

$$\lim_{r \rightarrow 0} \frac{\text{Vol}(B_r(x) \cap A)}{\text{Vol}(B_r(x))} = 1,$$

where Vol denotes the n -dimensional Lebesgue-measure.

References

- [1] C. Borgs, J. Chayes, L. Lovasz, V. T. Sós, B. Szegedy and K. Vesztergombi *Graph limits and parameter testing*. STOC'06: Proceedings of the 38th Annual ACM Symposium on Theory of Computing, 261–270, ACM, New York, 2006.
- [2] T. Gowers, *Quasirandomness, counting and regularity for 3-uniform hypergraphs*. Combin. Probab. Comput. **15** (2006), no. 1-2, 143–184.
- [3] P. R. Halmos, *Measure Theory* Van Nostrand Company, Inc., New York, N. Y., 1950.
- [4] Y. Ishigami, *A Simple Regularization of Hypergraphs*
preprint <http://arxiv.org/abs/math/0612838>
- [5] P. E. Loeb, *Conversion from nonstandard to standard measure spaces and applications in probability theory*. Trans. Amer. Math. Soc. **211** (1975), 113–122.
- [6] L. Lovasz, B. Szegedy, *Limits of dense graph sequences*. J. Combin. Theory Ser. B **96** (2006), no. 6, 933–957.
- [7] D. Maharam, *On homogeneous measure algebras*. Proc. Nat. Acad. Sci. U. S. A. **28**, (1942). 108–111.
- [8] B. Nagle, V. Rödl and M. Schacht, *The counting lemma for regular k -uniform hypergraphs*. Random Structures Algorithms **28** (2006), no. 2, 113–179.
- [9] V. Rödl, J. Skokan, *Regularity lemma for k -uniform hypergraphs*. Random Structures Algorithms **25** (2004), no. 1, 1–42.
- [10] J. Solymosi, *A note on a question of Erdős and Graham*. Combin. Probab. Comput. **13** (2004), no. 2, 263–267.
- [11] T. Tao, *A variant of the hypergraph removal lemma*. J. Combin. Theory Ser. A **113** (2006), no. 7, 1257–1280.

Gábor Elek Alfred Renyi Institute of the Hungarian Academy of Sciences POB 127, H-1364, Budapest, Hungary, elek@renyi.hu

Balázs Szegedy University of Toronto, Department of Mathematics, St George St. 40, Toronto, ON, M5R 2E4, Canada